



34110 9th Ave S, Federal Way, WA 98003 | [www.fikes.com](http://www.fikes.com) | 253.854.2439

---

## Stinging Insect Treatment Prep Sheet

(Paper Wasp, Yellow Jackets, Carpenter Bees, Hornets, Mud Daubers, etc.)

In order for our company to provide the most effective service possible, we ask that before our arrival you review and complete all the necessary preparations listed below. **Please be prepared to leave the area for up to 4 hours.** If you notice wet product, please refrain from entering this area until it has dried. The following steps will ensure that your Technician is able to provide an effective and safe treatment.

Fikes reserves the right to deny treatment if property is improperly prepared according to these instructions. Treatment charges may still be applied.

### How To Prepare For The Service:

Please take the following steps prior to our service to ensure that we can treat the most spaces during our time:

- Ensure perimeter of home that is in proximity of the nest is accessible by moving items away from the foundation of the home by 24 inches. ie. planters, patio furniture, etc.
- Never seal over the entry point of where wasps have been seen going into brick or stonework.
- Remove children, pets, children's toys, pet toys, pet food, bird food, and other exposed food from the effected area.
- Close all windows.

### What You Can Expect From The Treatment:

The treatment will remain effective only if the following steps are taken after your home is serviced:

- DO NOT use other pest control chemicals (powders or sprays) or cleaning products anywhere where treatment was applied (treatment will be applied on foundation of home, eaves, under and around decks, around exterior structures such as sheds, etc.).
- You may place back all exterior furniture, toys, and other items after product is dry, 2-4 hours after treatment.
- Do not water near the treatment for at least 24 hours after treatment as this will break down what has been applied and reduce its effectiveness.
- Report back if you are still experiencing concerns 2 weeks after treatment.